SPECIMEN MATERIAL v1.1

Please write clearly in	block capitals.		
Centre number		Candidate number	
Surname			
Forename(s)			
Candidate signature			

A-level CHEMISTRY

Paper 1: Inorganic and Physical Chemistry

Specimen materials (set 2)

Materials

For this paper you must have:

- the Periodic Table/Data Booklet, provided as an insert (enclosed)
- a ruler with millimetre measurements
- a scientific calculator, which you are expected to use where appropriate.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of the page.
- Answer all questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 105.

Time allowed: 2 hours

For Examiner's Use		
Question	Mark	
1		
2		
3		
4		
5		
6		
7		
8		
9		
10		
11		
TOTAL		

		04
	Answer all questions in the spaces provided.	
0 1	This question is about the chemistry of some Group 2 elements.	
01.1	Write an equation for the reaction of calcium with water at 25 °C and predict a	
	[2 ma	rks]
	Equation	
	рН	
0 1 . 2	State the trend in solubility, in water, of the Group 2 sulfates from magnesium to barium.	
	[1 m	ark]
0 1 . 3	State a reagent that can be used to test for the presence of sulfate ions and write	а
	[2 ma	rks]
	Reagent	
	Equation	
0 1 . 4	Explain why the melting point of calcium sulfate is high.	
	[2 ma	rks]

02	The following tests were carried out to identify an unknown green salt ${f Y}$.
	An aqueous solution of Y gave a cream precipitate of compound A when reacted with silver nitrate solution.
	Compound A gave a colourless solution when reacted with concentrated ammonia solution.
	Another aqueous solution of Y gave a green precipitate B when reacted with sodium carbonate solution.
	The green precipitate B was filtered and dried and then reacted with sulfuric acid to give a pale green solution containing compound C and a colourless gas D .
02.1	Identify by name or formula the compounds A , B , C , D and Y . [5 marks]
	Identity of A
	Identity of B
	Identity of C
	Identity of D
	Identity of Y
02.2	Write the simplest ionic equation for the reaction of silver nitrate solution with the anion that is present in compound Y . [1 mark]
02.3	Write the simplest ionic equation for the reaction that occurs between the green
	[1 mark]



		_
03.3	State why iodine does not conduct electricity.	[1 mark]
0 3 . 4	Deduce an equation for the formation of hydrogen iodide from its elements.	[1 mark]
	Question 3 continues on the next name	
	Question 5 continues on the next page	

03.5	The triiodide ion is formed when an iodine molecule is bonded to an iodide io	on.
	What is the formula of ammonium triiodide?	
	Tick (✓) one box.	[1 mark]
	NH ₃ I ₃	ני וומיאן
	NH ₃ I ₄	
	NH ₄ I	
	NH ₄ I ₃	
03.6	Draw the shape of the IF ₃ molecule and the shape of the IF ₄ ^{$-$} ion. Include any lone pairs of electrons that influence each shape.	[2 marks]
03.7	Deduce the oxidation state of iodine in the following species.	[2 marks]
	Ba(IO ₃) ₂	
	[H ₄ IO ₆] ⁻	

Г

0 4	Iron forms many complexes that contain iron in oxidation states +2 and +3.
04.1	Hexaaquairon(III) ions react with an excess of hydrochloric acid in a ligand substitution reaction.
	Write an equation for this reaction. [1 mark]
04.2	Explain why the initial and final iron(III) complexes in the equation in Question 04.1 have different shapes. [2 marks]
04.3	Hexaaquairon(II) ions react with an excess of $H_2NCH_2CH_2NH_2$ in a ligand substitution reaction. Draw the structure of the iron(II) complex formed showing its charge. [2 marks]
	Question 4 continues on the next page

04.4	Hexaaquairon(II) ions react with an excess of $H_2NCH_2CH_2NH_2$ in a ligand substitution reaction.
	Which of the following shows the correct change in entropy for a reaction of hexaaquairon(II) ions with $H_2NCH_2CH_2NH_2$?
	Tick (✓) one box. [1 mark]
	change in entropy is negative
	change in entropy is close to zero
	change in entropy is positive
04.5	The percentage of iron(II) sulfate in iron tablets can be determined by titration with potassium manganate(VII) in acidic solution.
	Deduce an ionic equation for the reaction of iron(II) ions with manganate(VII) ions. [1 mark]

04.6	A student dissolved 1980 mg of iron tablets in an excess of dilute sulfurior. The solution was titrated with 0.0200 mol dm ⁻³ potassium manganate(VII A 32.50 cm ³ volume of potassium manganate(VII) solution was required end point in the titration.	c acid. II) solution. to reach the
	Calculate the percentage of iron in the sample of iron tablets. Give your answer to the appropriate number of significant figures.	[4 marks]
	Percentage	%
04.7	State the colour change at the end point in this titration.	[1 mark]



0 5 . 3	Suggest how the enthalpy of lattice formation of NaCl compares with that of NaF Justify your answer. [3 marks]
	How enthalpies of formation compare
	Justification
0 5 . 4	Calculate the volume, in cm ³ , of fluorine gas at 298 K and 100 kPa required to produce 1.00 g of sodium fluoride by reaction with an excess of sodium.
	The gas constant $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ [4 marks]
	Volume cm ³

06	Table 2 s	hows some standard ele	ectrode potential	l data.	
			Table 2		
		Electrode half	-equation	<i>E</i> ^e / V	
		Cl ₂ (g) + 2e ⁻ _	→ 2Cl ⁻ (aq)	+1.36	
		Ag⁺(aq) + e⁻ _	→ Ag(s)	+0.80	
		Fe ³⁺ (aq) + e ⁻ –	→ Fe ²⁺ (aq)	+0.77	
		Sn ²⁺ (aq) + 2e [−] –	→ Sn(s)	-0.14	
		Fe ²⁺ (aq) + 2e ⁻ –	\rightarrow Fe(s)	-0.44	
06.1	Use data	a from Table 2 to deduc	e the species the tation for the cel	at is the best of	xidising agent. [1 mark] sure the standard [2 marks]
06.3	A cell wa consiste in a solu Calculat the nega	as made by connecting f d of silver in a solution o tion of tin(II) nitrate solu e the EMF of this cell ar ative electrode.	two half-cells wit of silver nitrate so tion. nd write a half-ec	th a salt bridge olution and the quation for the	 One half-cell other consisted of tin reaction that occurs at [2 marks]
	EMF Half-equ	ation			



0 7	Hydrogen can be manufactured by the reaction of methane with steam. An equilibrium is established as shown by the equation.
	$CH_4(g) + H_2O(g) \rightleftharpoons CO(g) + 3H_2(g)$
07.1	Use Le Chatelier's principle to predict the effect on the equilibrium yield of hydrogen if the overall pressure is increased. Explain your answer. [3 marks]
	Effect on yield
07.2	Explain why the equilibrium yield of hydrogen is unchanged if a catalyst is used in the reaction. [2 marks]

0 7 . 3 Table 3 shows the standard enthalpy of formation and the standard entropy for each substance in this equilibrium reaction. Table 3 $CH_4(g)$ $H_2O(g)$ CO(g) $H_2(g)$ $\Delta_{f}H^{\Theta}$ / kJ mol⁻¹ -75 -242 -111 0 S^e /J K⁻¹ mol⁻¹ 186 189 198 131 Use data from Table 3 to calculate the standard enthalpy change for this equilibrium reaction. [2 marks] kJ mol⁻¹ Standard enthalpy change 0 7 . 4 Use your answer from Question 07.3 and the entropy data from Table 3 to calculate the minimum temperature, in °C, needed for this reaction to be feasible. Give your answer to the appropriate number of significant figures. (If you did not complete Question 07.3 you should assume a value of 120 kJ mol⁻¹ for the standard enthalpy change. This is not the correct value). [5 marks] Minimum temperature °C

0 8 0 8 . 1	This question is about Brønsted–Lowry acids. Give the meaning of the term Brønsted–Lowry acid.	[1 mark]
08.2	What is meant by the term strong when describing an acid?	[1 mark]
08.3	At 298 K, 25.0 cm ³ of a solution of a strong monoprotic acid contained 1.45×10^{-3} mol of hydrogen ions. Calculate a value for the pH of this solution. Give your answer to 2 decimal places.	[2 marks]
	рН	









[2 marks]

The student was provided with samples of three different indicators.

to identify the most suitable indicator.

Suggest how the practical procedure in Question 09.2 could be refined by the student

09.3

Turn over for the next question

1 0	Acid X reacts with methanol to form an ester Y .
	$\begin{array}{c} CH_2COOH \\ \\ CH_2COOH \end{array} + 2CH_3OH \end{array} \rightleftharpoons \begin{array}{c} CH_2COOCH_3 \\ \\ CH_2COOCH_3 \end{array} + 2H_2O \qquad \Delta H^{\Theta} = -15 \text{ kJ mol}^{-1} \\ CH_2COOCH_3 \end{array}$
	X Y
10.1	Write an expression for the equilibrium constant, K_{c} , for this reaction. Use X and Y in your expression
	[1 mark]
10.2	A mixture of 0.32 mol of acid X and 0.84 mol of CH ₃ OH was allowed to reach equilibrium in the presence of a small amount of catalyst. The equilibrium mixture formed contained 0.26 mol of ester Y .
	Calculate the amounts, in moles, of X , CH ₃ OH and H ₂ O in this equilibrium
	mixture.
	Amount of X
	Amount of CH ₃ OH
	Amount of H ₂ O
10.3	Calculate the value of <i>K</i> _c and state the units. [3 marks]
	K _c units
10.4	Predict the effect on K_c if the reaction is carried out at a lower temperature. [1 mark]



1 1 .

A sample of the element Q consists of several isotopes. All of the Q⁺ ions in the sample of Q that has been ionised in a TOF mass spectrometer have the same kinetic energy.

kinetic energy of each ion = $\frac{1}{2}mv^2$

where *m* is the mass, in kg, of one ion of an isotope and *v* is the velocity of an ion in m s⁻¹

 $v = \frac{d}{t}$

where d is the length, in m, of the flight tube and t is the time taken, in s, for an ion to reach the detector

The time of flight of a ${}^{82}\mathbf{Q}^{+}$ ion is 1.243 × 10⁻⁵ s.

Calculate the time of flight of the ${}^{86}\mathbf{Q}^+$ ion.

[3 marks]

_ S

6

Time of flight of the ${}^{86}\mathbf{Q}^+$ ion _____

END OF QUESTIONS